

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/373454460>

# Diversity and species composition of microbiota associated with dengue mosquito breeding habitats: A cross-sectional study from selected areas in Udapalatha MOH division, Sri Lanka

Article in *Asian Pacific Journal of Tropical Medicine* · August 2023

DOI: 10.4103/1995-7645.380722

CITATIONS

0

READS

68

3 authors:



**Yasoda Kumari**

University of Kelaniya

3 PUBLICATIONS 0 CITATIONS

SEE PROFILE



**Deepika Amarasinghe**

University of Kelaniya

97 PUBLICATIONS 414 CITATIONS

SEE PROFILE




**Dr Koshila Ranasinghe**

University of Kelaniya

30 PUBLICATIONS 55 CITATIONS

SEE PROFILE



## Original Article

## Asian Pacific Journal of Tropical Medicine

apjtm.org



doi: 10.4103/1995–7645.380722

Impact Factor: 3.1

## Diversity and species composition of microbiota associated with dengue mosquito breeding habitats: A cross-sectional study from selected areas in Udapalatha MOH division, Sri Lanka

Yashoda Kumari, Deepika Amarasinghe<sup>✉</sup>, Koshila Ranasinghe

Department of Zoology and Environmental Management, Faculty of Science, University of Kelaniya, Dalugama, Kelaniya GQ 11600, Sri Lanka

### ABSTRACT

**Objective:** To determine the diversity of microbiota associated with different breeding habitats of dengue vector mosquitoes *Aedes (Ae.) aegypti* and *Ae. albopictus* and to identify any parasitic, epibiont, pathogenic, competitive or predatory species.

**Methods:** Sampling was performed from a variety of breeding habitats using dipping, pipetting and siphoning techniques. Microbiota in water samples were preserved using Rose Bengal solution and Lugol's iodine, and were identified. Live samples of microbiota were kept under laboratory conditions to observe any pathogenic or parasitic microbiota interacting with larvae.

**Results:** A total of eleven microbiota species (*Canthocamptus staphylinus*, *Canthocamptus microstaphylinus*, *Parastenocaris brevipes*, *Lepadella ovalis*, *Lepadella patella*, *Rotatoria rotatoria*, *Rotatoria macrura*, *Asplanchna brightwelli*, *Trichocerca rattus*, *Euglena variabilis*, and *Flagilaria capucina*) belonging to four (4) phyla (Arthropoda, Rotifera, Euglenozoa, and Ochrophyta) and 8 microbiota species belonged to four phyla (Arthropoda, Rotifera, Euglenozoa, and Ochrophyta) were identified from *Ae. aegypti* and *Ae. albopictus* breeding habitats respectively. There was a higher percentage (54.54%) of larval habitats positive for the secondary vector *Ae. albopictus* than through the primary vector *Ae. aegypti* in the Gampola urban area indicating higher possibility of transmitting the dengue virus through the secondary vector. However, no pathogenic or parasitic ciliates on mosquito larvae were encountered in the present study. Those findings may be due to sampling mainly from temporary container-type breeding habitats.

**Conclusions:** The relative distribution of microbiota associated with mosquito species differed significantly among *Ae. aegypti* and *Ae. albopictus*. The overall findings of this study could help in implementing novel eco-friendly vector-control strategies in the study area.

**KEYWORDS:** *Aedes*; Biological; Mosquito-control; Vectors

### 1. Introduction

In both tropical and temperate climates, mosquitoes spread a variety of vector-borne diseases to people, making them a significant group of pathogen-transmitting vectors[1]. Therefore, the study of mosquitoes' ecological and environmental factors that determine their abundance is a vital necessity[2,3]. The selectivity of a suitable oviposition site is an important factor in determining the success of

#### Significance

Identification of parasitic, epibiont, pathogenic, competitive or predatory microbiota in larval habitats and their interactions with associated mosquito larvae, in terms of controlling agents, would be beneficial for potential larval-controlling approaches. The degree of such parasitic, pathogenic, or predatory effects may vary with the geographical location. During the present study, a total of eleven and eight microbiota species were identified from *Aedes aegypti* and *Aedes albopictus* breeding habitats respectively from Udapalatha MOH division. The relative distribution of microbiota associated with mosquito species differed significantly among the *Aedes aegypti* and *Aedes albopictus* revealing the relationship of microbiota abundance with different mosquito species which helps in implementing novel vector-control strategies in the study area in an eco-friendly manner.

<sup>✉</sup>To whom correspondence may be addressed. E-mail: deepika@kln.ac.lk

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**For reprints contact:** reprints@medknow.com

©2023 Asian Pacific Journal of Tropical Medicine Produced by Wolters Kluwer-Medknow.

**How to cite this article:** Kumari Y, Amarasinghe D, Ranasinghe K. Diversity and species composition of microbiota associated with dengue mosquito breeding habitats: A cross-sectional study from selected areas in Udapalatha MOH division, Sri Lanka. Asian Pac J Trop Med 2023; 16(8): 363-370.

**Article history:** Received 5 February 2023  
Accepted 14 August 2023

Revision 2 August 2022  
Available online 28 August 2023

the life cycle. Adult mosquitoes use different cues like visual, tactile, etc. to select oviposition locations[4]. The factors that influence oviposition site selectivity mainly include the water quality of the breeding habitat. It is one of the most critical parameters that determine the success of egg hatching and the development of progeny. Therefore, females choose breeding places based on biotic and abiotic constituents of water[5,6]. Such abiotic parameters include pH, salinity, breeding site temperature, ionic concentrations and vegetation[7]. Meanwhile, biotic parameters also include competitors, predators, and the presence of parasites which influence mosquito larval development[8–10].


Naturally occurring microbiota is another biotic factor associated with mosquito breeding habitats. Among this diverse microbiota, food sources for mosquitoes, competitors, parasites, epibionts and predators may present. Such parasitic or pathogenic microbiota species may cause lethal effects on mosquito larvae[11]. Further, there are competitors such as bacteria, protists, and algae that consume the same food items as mosquito larvae. Predatory microbes such as *Calanoida*, *Harpacticoida*, and *Cladocera* could interfere with the development of mosquito larvae, thereby influencing their survival rates[12]. Meanwhile, it may negatively affect the egg-laying of gravid female mosquitoes[10]. Therefore, some microbiota in the mosquito breeding habitats may operate as natural mosquito larvae biocontrol agents[11].

Mosquitoes harbor communities of symbiotic microbes exhibiting various functions in their digestive tract including promoting or assisting the gut infection of incoming pathogens and significantly contributing to disease transmission and host-parasite interactions. The naturally occurring microbiota acquired by larvae from breeding habitats may establish as symbiotic flora in mosquito larval gut. Therefore associated microbiota species can affect the ability of mosquitoes to transmit disease-causing pathogens too[13]. Identification of naturally occurring microbiota and their interactions with mosquito larvae, in terms of epibiont, parasitic, pathogenic, competitive or predatory organisms against mosquito larvae as controlling agents would be beneficial for potential larval controlling approaches in an environmental-friendly manner.

## 2. Materials and methods

### 2.1. Study area

Gampola ( $7^{\circ} 7' 36.3972''$  N,  $80^{\circ} 33' 52.8372''$  E) is a town located in Kandy District, Central Province, Sri Lanka, and consists of an extent of land of about 94.02 km<sup>2</sup>. From Udapalatha MOH division, Gampola urban area was selected as the study area for the present study.



**Figure 1.** Sampling locations from the selected study site; Udapalatha MOH Division.

### 2.2. Sampling of mosquito breeding habitats for microbiota and mosquito larvae

Sampling was performed bi-monthly from November 2021 to January 2022. A total of 50 breeding habitat with mosquito larvae was collected from the sampling site, including blocked drains, discarded pots and plastic cups accumulated with rain water, discarded coconut shells, bamboo tree holes, ornament ponds, leaf axils, discarded tires and discarded roof tiles.

Breeding sites were selected within the district randomly, and each sampling site was geo-referenced (GARMIN-etrex SUMMIT) (Figure 1). Breeding habitats were categorized as man-made and natural[14]. Water samples were collected using a standard 250 mL dipper. When dipping is impossible, sampling was performed using pipetting or siphoning methods (maximum 250 mL) into a larval-rearing container (Height 12 cm, Diameter 6.5 cm). Dipping was performed for larger temporary habitats with a greater volume of water. For this, the metal scooper (250 mL volume scoop with a 30 cm long handle) was held vertically into the water body and a sample of water was taken maximum at the handle depth to comprise subsurface and bottom layers. When dipping is impossible, in small and flatwater sources, sampling was performed by pipetting out the

water using a pasture pipette; siphoning was done in places such as tree holes, leaf axils, and tires where both dipping and pipetting is impossible.

Both mosquito and microbiota sampling was performed from each habitat. At the sampling time, a collected water sample from the breeding habitat was divided into three plastic containers (6.5 cm width, 12 cm height); 2 for identification of microbiota and 1 for mosquito larval species identification and observing interactions of microbiota with live mosquito larvae. Two of them were immediately preserved in two methods using Rose Bengal stain (5% formalin with 0.04% Rose Bengal stain) solution and 5% Lugol's solution for microbiota identification. The remaining sample was kept as it is (non-preserved) and covered with a small-sized mesh net for getting live observations. Mosquito species identification was performed using standard identification keys [7,15,16]. All samples were labeled and transferred carefully into the laboratory for further processing.

### 2.3. Identification of microbiota

A total of 1.0 mL aliquot of preserved sample was examined under the compound microscope ( $\times 100$  magnification) (Olympus  $\times C21$ , Japan) using a Sedgwick rafter (S-R) cell (50 mm length, 20 mm width, 1 mm deep) and HYDRO-BIOS phytoplankton chamber (dimensions: 33 mm  $\times$  33 mm; thickness: 1 mL) for quantifying the microbiota. The sample was well shaken before taking the aliquot for observation. Microbiota species/taxa were identified at taxa/species level using temporary slide mounts. Microbiota were identified using standard identification keys ( $\times 400$  magnification) [17–19]. The non-preserved sample was observed daily until the pupation of mosquito larvae there.

### 2.4. Larval rearing and taxonomic identification

The mosquito larvae were first separated to the genus level and classified into instar stages ( I & II and III or IV). The larvae were identified under a Binocular light microscope (Olympus C21, Japan) using morphological taxonomic keys [7,15,16,20,21].

### 2.5. Determination of possible parasitic/pathogenic/epibiont or predatory microbiota against mosquito larvae

Regular daily observations were made, from non-preserved water samples for any significant survival change or reduction of mosquito larval count or any change of motility of live mosquito larvae due to biologically affecting microbiota species/taxa associated. Epibiont/symbiont or predator if any, was observed by micro pipetting 1 mL sample into Sedgwick rafter (S-R) cell and observing under the microscope ( $\times 40$  and  $\times 100$  magnifications).

### 2.6. Data analysis

Occurrence frequencies of microbiota species were categorized as constant for species found in more than 50% of the collections; common when found between 25% and 50% of the collections; and accidental or rare species when found in less than 25% of the collections [22]. Microbiota alpha diversity ( $\alpha$ ) was calculated for each breeding habitat type as the total number of species in the sampling periods, and  $\alpha$  medium was calculated as the average between the diversity for the system of the same type; gamma ( $\gamma$ ) diversity was estimated using the total number of species from all samples.

Beta diversity ( $\beta$ ) was estimated by measuring the species turnover using the  $\beta$ -1 index [23], which measures the amount that

**Table 1.** List of breeding habitats positive for *Aedes aegypti* and *Aedes albopictus* mosquito immature stages encountered from selected area in Udapalatha MOH Division, Gampola, Sri Lanka.

Breeding habitat	Type of breeding habitat	<i>Aedes aegypti</i>	<i>Aedes albopictus</i>
Plastic container (11)	Man-made (11)	+	+
Metal container (14)	Man-made (14)	+	+
Concrete slabs (2)	Man-made (2)	+	+
Glassware (1)	Man-made (1)	+	-
Tires (2)	Man-made (2)	+	-
Leaf axils (5)	Natural (5)	+	+
Tree holes (1)	Natural (1)	+	-
Coconut shells (2)	Man-made (2)	+	+
Bamboo tree (2)	Natural (2)	+	+
Ornamental pond (1)	Man-made (1)	+	-
Discarded roof tile (1)	Man-made (1)	-	+
Clay pots (2)	Man-made (2)	-	+

Note: The number of habitats sampled is included in parenthesis.

regional diversity exceeds mean alpha diversity. It was calculated by the formula  $\beta-1 = [(S/\alpha_{\text{mean}})-1]/[N-1] \times 100$ , where  $S$  is the regional diversity or total richness (the number of species per each sampling site);  $\alpha_{\text{mean}}$  is the mean  $\alpha$  diversity (mean number of species) for each site in each period;  $N$  is the number of sites of the period. Beta-diversity over 50% indicates high heterogeneity in microbiota composition among systems; between 20% and 50% indicates intermediate heterogeneity; and below 20% indicates low heterogeneity[23]. The microbiota species diversity was also estimated according to the indices of Shannon and Wiener[24] and evenness[25].

The Chi-square test of independence was used to evaluate the significance of the distribution of different microbiota species among different breeding sites of *Aedes (Ae.) aegypti* and *Ae. albopictus* in the study area.

### 3. Results

#### 3.1. Habitat positivity

A total of 44 breeding habitats were observed with the presence of *Ae. aegypti* and *Ae. albopictus* larvae from the total sampled 50 breeding habitats. Moreover, a total of 12 temporary key breeding sites were identified, with *Aedes* larvae (Table 1). Leaf axils, tree holes and bamboo trees were found in the study area as natural mosquito breeding habitats. The majority of the sampled breeding habitats belonged to the category of man-made temporary micro-breeding habitats (Table 1). Such temporary micro-breeding habitat types were positive for *Ae. aegypti* and *Ae. albopictus* mosquito immature stages. The highest mosquito larval diversity and abundance were found in metal containers. Plastic containers, metal containers, concrete slabs, leaf axils, coconut shells, and bamboo trees were found positive for both *Ae. aegypti* and *Ae. albopictus*. Besides, glassware, tires, ornamental ponds, and tree holes were positive for *Ae. aegypti*, and discarded roof tiles and clay pots were positive for *Ae. albopictus* (Table 1).

*Ae. albopictus* showed a relatively higher distribution and abundance over *Ae. aegypti*. No co-existing of *Ae. aegypti* and *Ae. albopictus* was found in the samples during the present study.

#### 3.2. Diversity and occurrence of microbiota from different mosquito breeding habitats of *Ae. aegypti* and *Ae. albopictus*

A total of eleven (11) microbiota species belonging to four (4) phyla (Arthropoda, Rotifera, Euglenozoa, and Ochrophyta) were identified from 20 different mosquito breeding habitats of *Ae. aegypti*; while 8 microbiota species belonging to four phyla (Arthropoda, Rotifera, Euglenozoa, and Ochrophyta) were identified from 20 breeding habitats of *Ae. albopictus* (Figure 2).




Figure 2. Percentage abundance of microbiota phyla encountered from different mosquito breeding habitats of *Aedes aegypti* and *Aedes albopictus*.




Figure 3. Occurrence of microbiota phyla encountered from mosquito breeding habitats. (A) *Aedes aegypti* (B) *Aedes albopictus*.

The phylum Rotifera gave the highest percentage abundance (51.66%) of total microbiota and the phylum Euglenozoa gave the lowest percentage abundance (1.02%) for *Ae. aegypti* breeding habitats. Meanwhile for *Ae. albopictus* breeding habitats, phylum Arthropoda showed the highest percentage of abundance (40.64%), followed by phylum Rotifera (40.27%) and the phylum Ochrophyta had the lowest percentage of abundance (6.72%) (Figure 2).

Mosquito breeding habitat types occupied with *Ae. aegypti* larvae, exhibited a diversity of microbiota belonging to four phyla, with a higher abundance of phylum Rotifera in plastic containers and glassware. Rotifers exhibited a wide range of morphological variations within breeding habitats. *Rotaria rotatoria* had the highest abundance of the total rotifers found from *Ae. aegypti* breeding habitats (Supplementary Table 1). The abundance of the phylum

**Table 2.** Evenness, Shannon diversity, alpha ( $\alpha$ ) medium, beta ( $\beta$ ), and gamma ( $\gamma$ ) diversities of type of habitats.

Breeding habitat	Mosquito species	No. of habitats positive for larvae	Alpha ( $\alpha$ ) medium	Beta ( $\beta$ )	Gamma ( $\gamma$ )	Shannon-Weiner diversity	Evenness
Plastic container (n=11)	<i>Aedes aegypti</i>	5	2	0.5	6	1.3	0.7
	<i>Aedes albopictus</i>	6	1	0.8	5	1.3	0.8
Concrete slabs (n=2)	<i>Aedes aegypti</i>	1	1	0.0	1	0.0	0.0
	<i>Aedes albopictus</i>	1	2	0.0	2	0.7	1.0
Metal container (n=14)	<i>Aedes aegypti</i>	4	2	0.3	4	1.2	0.9
	<i>Aedes albopictus</i>	10	2	0.2	6	1.4	0.8
Leaf axils (n=5)	<i>Aedes aegypti</i>	3	3	0.7	7	1.7	0.9
	<i>Aedes albopictus</i>	2	2	0.5	3	1.1	1.0
Bamboo tree (n=2)	<i>Aedes aegypti</i>	1	3	0.0	3	1.1	1.0
	<i>Aedes albopictus</i>	1	1	0.0	1	0.0	0.0
Coconut shells (n=2)	<i>Aedes aegypti</i>	1	1	0.0	1	0.0	0.0
	<i>Aedes albopictus</i>	1	1	0.0	1	0.0	0.0
Glassware (n=1)	<i>Aedes aegypti</i>	1	2	0.0	2	0.5	0.8
Tires (n=2)	<i>Aedes aegypti</i>	2	1	1.0	2	0.4	0.6
Ornamental pond (n=1)	<i>Aedes aegypti</i>	1	1	0.0	1	0.0	0.0
Tree holes (n=1)	<i>Aedes aegypti</i>	1	1	0.0	1	0.0	0.0
Discarded roof tile (n=1)	<i>Aedes albopictus</i>	1	2	0.0	2	0.7	0.9
Clay pots (n=2)	<i>Aedes albopictus</i>	2	1	0.5	3	1.0	0.9

Arthropoda was found to be dominant in metal containers and leaf axils (Figure 3).

Considering the breeding habitats occupied by *Ae. albopictus*, phylum Arthropoda showed the highest abundance. Phylum Arthropoda was dominant in plastic containers and leaf axils. Further, metal containers were identified as the breeding habitat with the highest number of microbiota species occurrence with *Ae. albopictus* larvae (Figure 3). Phylum Rotifera was found as dominant in metal containers. Discarded roof tiles exhibit the highest abundance of microbiota from the phylum Euglenozoa. Bamboo trees, coconut shells and concrete slabs had the presence of only one microbiota phyla; Arthropoda, Ochrophyta, and Rotifera, respectively (Figure 3).

### 3.3. Occurrence frequencies of microbiota species in different types of breeding habitats of *Ae. aegypti* and *Ae. albopictus*

*Canthocamptus staphylinus* has existed as a common microbiota species in many breeding habitat types such as concrete slab (100.00%), plastic containers (50.03%), associating with *Ae. aegypti* larvae and in bamboo trees (100.00%) and clay pot (58.83%) associating with *Ae. albopictus* larvae (Supplementary Table 1). Interestingly it has existed as an accidental and rare species also in other breeding habitat types. Therefore, *Canthocamptus staphylinus* has been identified as a species that shows all three possible occurrences in different types of breeding habitats (Supplementary Table 1).

The only microbiota species recorded from the study area that belonged to phylum Ochrophyta was *Euglena capucina* and it was

encountered in plastic containers, tires, leaf axils and coconut shells with *Aedes* larvae (Supplementary Table 1). *Ae. albopictus* larvae were more frequently occupied with two common microbiota species: *Canthocamptus staphylinus* (42.15%) and *Rotaria rotatoria* (32.35%) in metal containers. All other microbiota species had a rare occurrence in metal containers (Supplementary Table 1).

Considering the microbiota abundance in plastic containers associated with *Ae. aegypti*, *Rotaria rotatoria* was recognized as a constant species with a 51.73% occurrence, and all other microbiota recorded belonged to common or rare categories (Supplementary Table 1).

The highest Shannon Weiner diversity index and species richness/gamma ( $\gamma$ ) diversity of microbiota associated with *Ae. aegypti* larvae were recorded from leaf axils, while it was from metal containers for *Ae. albopictus* (Table 2). Further, the highest heterogeneity of microbiota associated with *Ae. albopictus* was recorded from plastic containers as it gave the highest beta diversity value of 0.8. Tires for *Ae. aegypti* had the highest beta ( $\beta$ ) diversity value of 1 (Table 2). Therefore, tires for *Ae. aegypti* and plastic containers for *Ae. albopictus* indicated a higher heterogeneity of microbiota composition which is associated with their biology.

Moreover, the plastic containers and metal containers for *Ae. aegypti* had beta ( $\beta$ ) diversity between 0.2 and 0.5, indicating intermediate heterogeneity of microbiota composition among the systems (Table 2). This observation corresponded to metal container, leaf axil, and clay pot breeding habitats for larvae of *Ae. albopictus*. The rest of the habitats were with a  $\beta$  diversity below 0.2, indicating low heterogeneity of microbiota within the breeding habitat (Table 2).

### 3.4. Variation of microbiota communities across mosquito species

The relative distribution of microbiota associated with mosquito species differed significantly among the *Ae. aegypti* and *Ae. albopictus* ( $\chi^2=486.091$ ;  $P<0.001$ ).

### 3.5. Identified microbiota parasitic/ pathogenic/ epibiont to mosquito larvae

Natural population of mosquitoes was kept under check by the activities of parasites/epibionts and pathogens on mosquito larvae, no such microbiota was identified from the present sampling site.

## 4. Discussion

The present study identified naturally occurring microbiota species associated with a variety of vector mosquito breeding habitats. The present study findings address the knowledge gap regarding information, recording a total number of 11 microbiota species from a variety of mosquito breeding habitats in the study area.

Although some vector-borne diseases like malaria and filariasis have been eliminated from Sri Lanka through successful control strategies, some vector-borne diseases, such as dengue, have increasing trends each year[26]. It has become a major public health and socio-economic concern in Sri Lanka. Two major vector mosquito species are responsible for dengue transmission in Sri Lanka; *Ae. aegypti* and *Ae. albopictus*. *Ae. aegypti* is known to be the primary vector, which is the predominant vector in urban areas, while *Ae. albopictus* is considered as the secondary vector which is the predominant vector in rural areas[27]. Thus, the present study considered microbiota associated with both species of *Aedes* mosquito breeding waters. *Ae. aegypti* were mostly encountered from plastic containers while *Ae. albopictus* were from metal containers. Containers that keep water for extended periods, such as artificial containers, build better or excellent mosquito breeding environments[28–31]. Present study findings revealed, both *Ae. aegypti* and *Ae. albopictus* were occupied in different types of breeding habitats, especially in temporary microhabitats, indicating the high risk of dengue vector distribution through increased accumulation of artificial containers. Further, there was a higher percentage (54.54%) of larval habitats positive for the secondary vector *Ae. albopictus* than through the primary vector *Ae. aegypti* in the Gampola urban area indicating higher possibility of transmitting the dengue virus through the secondary vector.

Previous larval surveys carried out in the Gampola study area also revealed that the *Ae. albopictus* was the most abundant species in the area, compared to *Ae. aegypti* [32].

Previous studies on microbiota inhabiting mosquito breeding habitats in different districts also reported a wide range of microbiota with some parasitic species. Ranasinghe *et al*[12] reported some pathogenic or parasitic ciliates, including *Vorticella microstoma*, *Zoothamnium* spp., and *Chilodinella* sp., from rice field habitats in Gampaha District. Cyanobacterial diet items for mosquito larvae like *Spirulina* (from plastic containers, tree holes, etc.), *Anabaena affinis* (from irrigation canals), *Scenedesmus armatus* (from drainage and tree holes) and *Scenedesmus bijuga* (from ponds, plastic containers, etc.) were also recorded[12].

Further, Ranasinghe *et al*[33] reported mosquito larval mortalities associated with high densities of *Vorticella* sp. and *Zoothamnium* sp. attached to the siphon and thoracic cuticular areas. Anyhow such pathogenic or parasitic ciliates were not encountered in the present study. A previous study carried out in the Kandy district recorded several algae species associated with both *Ae. aegypti* and *Ae. albopictus* larvae[12]. Most algae that inhabit breeding water serve as food sources, while some algal species like blue-green algae and cyanobacteria have lethal effects on mosquito larvae[34–36]. However, such algae species were not identified from the present study site. Those findings may be due to sampling from temporary container-type breeding habitats mainly, during the present study. For the development of microbiota in a breeding habitat, abiotic and other biotic factors associated play a key role. Further, there is comparatively less diversity of microbiota in a temporary container-type habitat, compared to the diversity recorded in literature from a natural habitat[12]. The present study confirms the above findings.

Recently conducted experiments have shown the adverse impacts of cladocerans such as competitors and cyclopoid copepods like predators on mosquito survival[37]. Competitors and predators are efficient as biocontrol agents against mosquito larvae[10,38, 39]. Antagonistic crustaceans such as *Mesocyclops aspericornis* and *Daphnia magna* cause the late development of mortality of early instar larvae of *Aedes* by predatory effects and interspecific competition for resources[3]. From the present study, the highest species richness of microbiota for *Ae. aegypti* was recorded from Phylum Rotifera including *Lepadella ovalis*, *Lepadella patella*, *Rotaria rotatoria*, *Rotaria* sp., *Asplanchna brightwelli* and *Trichocerca rattus* in a range of breeding habitats. Several prior studies have highlighted the significance of some rotifers against mosquito larvae survival[40]. The species *Asplanchna brightwelli* and *Trichocerca rattus* were competitors or predators on mosquito larvae[40]. The present study has not revealed such an association may be due to less abundance of rotifers in breeding waters in container habitats. Overall present study findings revealed that temporary-container breeding habitats harbor less diversity and abundance of naturally-occurring microbiota associated with mosquito larvae, compared to a natural breeding habitat.

A total of eleven microbiota species belonging to four phyla

(Arthropoda, Rotifera, Euglenozoa, and Ochrophyta) and 8 microbiota species belonged to four phyla (Arthropoda, Rotifera, Euglenozoa, and Ochrophyta) were identified from *Ae. aegypti* and *Ae. albopictus* breeding habitats respectively. The relative distribution of microbiota associated with mosquito species differed significantly among the *Ae. aegypti* and *Ae. albopictus*. The presence of dengue vectors in the study area in considerable numbers can cause public health concerns as dengue is one of the major challenges in these areas. Therefore, a study of this nature would be useful to identify the entomological potential for disease transmission and an update on microbiota associated with *Aedes* mosquito larvae would be facilitated for implementing appropriate future vector control interventions. Morphological identification of some microbiota that was too small, up to the species level served as a limitation of the study.

### Conflict of interest statement

We declare that there is no conflict of interest.

### Funding

The present research received funds from Department of Zoology and Environmental Management, University of Kelaniya, Sri Lanka.

### Acknowledgments

This work was supported by the Department of Zoology and Environmental Management, University of Kelaniya, Sri Lanka.

### Authors' contributions

Deepika Amarasinghe did the designing of the research, overall supervision and guidance of the research work and final editing and reviewing the manuscript. Yasoda Kumari performed sampling and data collection, data analysis and writing the manuscript. Koshila Ranasinghe did the supervision of the research work, data analysis, writing and reviewing of the manuscript. All authors read and approved the final manuscript.

### References

[1] Becker N, Zgomba M, Petric D, Dahl C, Boase C, Lane J, et al. Mosquitoes and their control. Springer Science & Business Media; 2003,

p. 453-485.

- [2] Simsek F. Seasonal larval and adult population dynamics and breeding habitat diversity of *Culex theileri* Theobald 1903 (Diptera: Culicidae) in the Golbasi district, Ankara. *Turk J Zool* 2004; **28**: 337-344.
- [3] Chaves L, Koenraadt CJM. Climate change and highland malaria: Fresh air for a hot debate. *Q Rev Biol* 2010; **85**(1): 27-55.
- [4] Day, J. Mosquito oviposition behavior and vector control. *Insects* 2016; **7**(4): 65.
- [5] Bentley MD, Day JF. Chemical ecology and behavioral aspects of mosquito oviposition. *Annu Rev Entomol* 1989; **34**(1): 401-421.
- [6] Barrera R, Amador M, Clark GG. Ecological factors influencing *Aedes aegypti* (Diptera: Culicidae) productivity in artificial containers in Salinas, Puerto Rico. *J Med Entomol* 2006; **43**(3): 484-492. doi: 10.1093/jmedent/43.3.484.
- [7] Amerasinghe F, Indrajith N, Ariyasena T. Physicochemical characteristics of mosquito breeding habitats in an irrigation development area in Sri Lanka. *Ceylon J Sci* 1995; **24**(2): 13-29.
- [8] Muturi EJ, Mwangangi J, Shililu J, Muriu S, Jacob B, Kabiru E, et al. Mosquito species succession and physicochemical factors affecting their abundance in rice fields in Mwea, Kenya. *J Med Entomol* 2007; **44**(2): 336-344.
- [9] Oyewole OI, Momol OO, Anyasor GN. Physicochemical characteristics of *Anopheles* breeding sites: Impact on fecundity and progeny development. *Afr J Environ Sci & Technol* 2009; **3**(12): 447-452.
- [10] Meyabeme Elono AL, Liess M, Duquesne S. Influence of competing and predatory invertebrate taxa on larval populations of mosquitoes in temporary ponds of wetland areas in Germany. *J Vector Ecol* 2010; **35**(2): 419-427.
- [11] Loria K. *Freshwater zooplankton communities as indicators of habitat quality: Testing responses*. Ph.D thesis submitted to University of Colorado at Boulder; 2017, p. 36.
- [12] Ranasinghe HAK, Amarasinghe LD. Naturally occurring microbiota associated with mosquito breeding habitats and potential parasitic species against mosquito larvae: A study from Gampaha District, Sri Lanka. *BioMed Res Int* 2020; 1-12. doi: 10.1155/2020/4602084.
- [13] Nilsson LKJ, de Oliveira MR, Marinotti O, Rocha EM, Håkansson S, Tadei WP, et al. Characterization of bacterial communities in breeding waters of *Anopheles darlingi* in Manaus in the Amazon basin malaria-endemic area. *Microb Ecol* 2019; **78**(4): 781-791.
- [14] Yee DA, Allgood D, Kneitel JM, Kuehn KA. Constitutive differences between natural and artificial container mosquito habitats: Vector communities, resources, microorganisms and habitat parameters. *J Med Entomol* 2012; **49**(3): 482-491.
- [15] Chelliah RV. Keys and illustrated key to the genera of mosquitoes of Sri Lanka (Diptera: Culicidae). *Contrib Am Entomol Inst* 1984; **7**(4): 1-84.
- [16] Rueda LM. Pictorial keys for the identification of mosquitoes (Diptera: Culicidae) associated with dengue virus transmission. *Zootaxa* 2004; **589**(1): 1-60.
- [17] Fernando CH, Weerawardhena SR. *A guide to the freshwater fauna of*



- Ceylon (Sri Lanka). Ceylon, Sri Lanka: Fisheries Research Station; 2002.
- [18]Abeywickrama BA, Abeywickrama L. *The genera of the freshwater algae of Sri Lanka*. Part 1. Colombo: National Science Council, Sri Lanka; 1979.
- [19]Corliss JO. *The ciliated protozoa: Characterization, classification, and guide to the literature*. 2nd ed. London, UK: Pergamon Press; 1979.
- [20]Gunathilaka N. Illustrated key to the adult female *Anopheles* (Diptera: Culicidae) mosquitoes of Sri Lanka. *Appl Entomol Zool* 2017; **52**(1): 69-77.
- [21]Gunathilaka N. Annotated checklist and review of the mosquito species (Diptera: Culicidae) in Sri Lanka. *J Insect Biodivers* 2018; **7**(3): 38-50.
- [22]Lobo E, Leighton G. Estructuras comunitarias de las fitocenosis planctónicas de los sistemas de desembocaduras de ríos y esteros de la zona central de Chile. *Rev Biol Mar Oceanogr* 1986; **22**: 1-29.
- [23]Harrison S, Ross SJ, Lawton JH. Beta diversity on geographic gradients in Britain. *J Anim Ecol* 1992; **62**: 151-158.
- [24]Shannon CE, Weaver W. *The mathematical theory of communication*. Urbana: The University of Illinois Press; 1949.
- [25]Pielou EC. *Ecological diversity*. New York: John Wiley; 1975.
- [26]Wijegunawardana NDAD, Gunawardene YINS, Chandrasena TGAN, Dassanayake RS, Udayanga NWBAL, Abeywickreme W. Evaluation of the effects of *Aedes* vector indices and climatic factors on dengue incidence in Gampaha District, Sri Lanka. *BioMed Res Int* 2019; 1-11. doi: 10.1155/2019/2950216.
- [27]Hawley W, Reiter P, Copeland R, Pumpuni C, Craig Jr. G. *Aedes albopictus* in North America: Probable introduction in used tires from Northern Asia. *Science* 1987; **236**(4805): 1114-1116.
- [28]Saleeza SNR, Rashid YN, Azirun MS. Mosquitoes larval breeding habitat in urban and suburban areas, Peninsular Malaysia. *IJABE* 2011; **5**: 599-603.
- [29]Philbert A, Ijumba JN. Preferred breeding habitats of *Aedes aegypti* (Diptera: Culicidae) mosquito and its public health implications in Dar es Salaam, Tanzania. *J Environ Res Manage* 2013; **4**(10): 344-351.
- [30]Wilson JJ, Sevarkodiyone SP. Spatial and temporal distribution of mosquitoes (Culicidae) in Virudhunagar district, Tamil Nadu, South India. *Int J Mosq Res* 2014; **1**(3): 4-9.
- [31]Sedaghat MM, Bozorg Omid F, Karimi M, Haghi S, Hanafi-Bojd AA. Modelling the probability of presence of *Aedes aegypti* and *Aedes albopictus* in Iran until 2070. *Asian Pac J Trop Med* 2023; **16**(1): 16-25.
- [32]Weeraratne TC, Perera BMD, Mansoor MACM, Karunaratne SHPP. Prevalence and breeding habitats of the dengue vectors *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) in the semi-urban areas of two different climatic zones in Sri Lanka. *Int J Trop Insect Sci* 2013; **33**(4): 216-226. doi: 10.1017/s174275841300026x.
- [33]Amarasinghe LD, Rathnayake ARLK. Prevalence of microfauna associated with different mosquito breeding habitats in a selected area of Sri Lanka. *Int J Curr Microbiol Appl Sci* 2014; **3**(5): 587-598.
- [34]Marten GG. Impact of the copepod *Mesocyclops leukarti pilosa* and green alga *Kirchneriella irregularis* upon larval *Aedes albopictus* (Diptera: Culicidae). *Bull Soc Vector Ecol* 1984; **9**(1): 1-5.
- [35]Marten GG. Mosquito control by plankton management: The potential of indigestible green algae. *J Trop Med Hyg* 1986; **89**: 213-222.
- [36]Ranasinghe HAK, Amarasinghe LD. Naturally occurring microbiota associated with mosquito breeding habitats and their effects on mosquito larvae. *BioMed Res Int* 2020; 1–11. doi: 10.1155/2020/4065315.
- [37]Thakur A, Kocher DK. Impact of antagonistic crustaceans on the population of *Aedes aegypti* L. larvae under laboratory conditions. *J Vector Borne Dis* 2020; **57**(1): 58-62. doi: 10.4103/0972-9062.308802.
- [38]Blaustein L, Chase JM. Interactions between mosquito larvae and species that share the same trophic level. *Ann Rev Entomol* 2007; **52**(1): 489-507. doi: 10.1146/annurev.ento.52.110405.091431.
- [39]Duquesne S, Kroeger I, Kutyniok M, Liess M. The potential of cladocerans as controphic competitors of the mosquito *Culex pipiens*. *J Med Entomol* 2011; **48**(3): 554-560. doi: 10.1603/me09282.
- [40]Duguma D, Kaufman MG, Simas Domingos AB. Aquatic microfauna alter larval food resources and affect development and biomass of West Nile and Saint Louis encephalitis vector *Culex nigripalpus* (Diptera: Culicidae). *Ecol Evolut* 2017; **7**(10): 3507-3519.

## Publisher's note

The Publisher of the *Journal* remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.