Sri Lanka Association for the Advancement of Science (SLAAS) Proceedings of the 56<sup>th</sup> Annual Session, Part 1 - Abstracts 27 November - 01 December, 2000. Colombo, Sri Lanka

D 263

E)

of

)CS

the

ya;

319 :lus

nly

itly osa

rere ysis nds

by file of

Effect of azadirachtin on midgut enzymes of cockroaches (Periplanata americana)

PA Paranagama<sup>1</sup>, KABCH Kodikara<sup>1</sup>, HMI Nishantha<sup>1</sup> and AM Mubarak<sup>2</sup>
<sup>1</sup>Department of Chemistry, University of Kelaniya, Kelaniya and <sup>2</sup>ITI, Colombo 7

Both *invivo* and *invitro* experiments were carried out to examine the effect of azadirachtin on midgut enzymes of adult female cockroaches. The activity of protease, invertase and anylase were measured separately in both azdirachtin treated cockroaches  $(3\mu g/g)$  body weight) and the control cockroaches using the methods described in Ishaaya et al,  $1971^2$  1974 3. The *invivo* results indicate that the azadirachtin treated cockroaches showed 50% reduction of the activity of all three enzymes and azadirachtin had no inhibitory effect on the activity of midgut enzymes *invitro*. The same experiment was carried out with the ligatured cockroaches in order to study the effect of neurohormones on the secretion of midgut enzymes. In contrast, similar results were obtained from the azadirachtin treated insects and the control insects suggesting the toxic effect of azadirachtin is associated with a disruption of endocrine events in cockroaches.

Financial assistance by NSF for research grant number RG/96/C/04 is acknowledged.